

REFERENCES

- Abas, F., Lajis, N.H., Israf, D.A., Khozirah, S. and Kalsom, Y.U. (2006). Antioxidant and nitric oxide inhibition activities of selected Malay traditional vegetables. *Food Chemistry*, 95, 566-573.
- Abdelwahab, S.I., Mohan, S., Elhassan, M.M., Al-Mekhlafi, N., Mariod, A.A., Abdul, A.B., Abdulla, M.A. and Alkharfy, K.M. (2011). Antiapoptotic and antioxidant properties of *Orthosiphon stamineus Benth* (Cat's Whiskers): Intervention in the Bcl-2-Mediated apoptotic pathway. *Evidence-Based Complementary and Alternative Medicine*, 2011, 1-11.
- Ahmadi, F., Kadivar, M., and Shahedi, M. (2007). Antioxidant activity of *Kelussia odoratissima* Mozaff. in model and food systems. *Food Chemistry*, 105, 57-64.
- Ak, T. and Gulcin, I. (2008). Antioxidant and radical scavenging properties of curcumin. *Chemico-Biological Interactions*, 174, 27-37.
- Alard, D., Wray, V., Grotjahn, L., Reznik, H. and Strack, D. (1985). Neobetanin: Isolation and identification from *Beta vulgaris*. *Phytochemistry*, 24 (10), 2383-2385.
- Almela, L., Mu noz, B.S., Lopez, J.A.F., Roca, M. and Rabe, V. (2006). Liquid chromatographic – mass spectrometric analysis of phenolics and free radical scavenging activity of rosemary extract from different raw material. *Journal of Chromatography A*, 1120, 221-229.
- Amarowicz, R., Pegg, R.B., Moghaddam, P.R., Barl, B., Well, J.A. (2004). Free-radical scavenging capacity and antioxidant activity of selected plant species from the Canadian prairies. *Food Chemistry*, 84(4), 551-562.
- Amez, B.N., Shigenaga, M.K., Hagen, T.M. (1993). Oxidants, antioxidants, and the degenerative disease of aging. *Proceedings of Natural Academy of Sciences. USA*, 90, 7915-7922.
- Atawodi, S.E. (2004). Antioxidant potential of african medicinal plants. *African Journal of Biotechnology*, 4(2), 128-133.
- Azerado, H.M.C. (2009). Betalains: properties, sources, applications, and stability – a review. *International Journal of Food Science and Technology*, 44, 2365-2376.
- Bailly, C. (2009). Ready for a comeback of natural products in oncology. *Biochemical Pharmacology*, 77, 1447-1457.
- Bandyopadhyay, M., Chakraborty, R., Raychaudhuri, U. (2008). Antioxidant activity of natural plant sources in dairy dessert (Sandesh) under thermal treatment. *Lebensmittel-Wissenschaft und-Technologie*, 41, 816-825.
- Barja, G. (2004). Free radicals and aging. *Trends in Neurosciences*, 27(10), 595-600.

- Bavec, M., Turinek, M., Grobelnik-Mlakar, S., Slatnar, A. and Bavec, F. (2010). Influence of Industrial and Alternative Farming Systems on Contents of Sugars, Organic Acids, Total Phenolic Content, and the Antioxidant Activity of Red Beet (*Beta vulgaris* L. ssp. *vulgaris* Rote Kugel). *Journal of Agricultural and Food Chemistry*, 58 (22), 11825–11831.
- Benzie, I.F. F. & Strain, J. J. (1996). The ferric reducing ability of plasma (FRAP) as a measure of antioxidant power: The FRAP assay. *Analytical Biochemistry*, 239, 70-76.
- Bin Li,H., Jiang, Y., Chun Wong, C., Wing Cheng, K., Feng Chen. (2007). Evaluation of two methods for the extraction of antioxidants from medicinal plants. *Analysis Bioanalysis Chemistry*, 388, 483-488.
- Bondet, V., Williams, W.B., Berset, C. (1997). Kinetics and mechanisms of antioxidant activity using the DPPH free radical method. *Lebensmittel-Wissenschaft und-Technologie*, 30(6), 609-615.
- Bor, M., Zdemir, F.O. and Turkan, I. (2003). The effect of salt stress on lipid peroxidation and antioxidants in leaves of sugar beet *Beta vulgaris* L. and wild beet *Beta maritima* L. *Plant Science*, 164, 77-84.
- Borenfreund E, Puerner J. (1984). *Journal of Tissue Culture Method* 9: 7-9.
- Brand-Williams, W., Cuvelier, M.E. and Berset, C. (1995). Use of a free radical method to evaluate antioxidant activity. *Lebensmittel-Wissenschaft und-Technologie/Food Science and Technology*, 28: 25-30.
- Bruno, G. (2008). Antioxidants against free radicals. *Huntington College of Health Sciences Literature Education Series On Dietary Supplements*.
- Buchner, F.L., Bueno-de-Mesquita, H.B., Ros, M.M., et al. (2010). Variety in fruit and vegetable consumption and the risk of lung cancer in the European prospective investigation into cancer and nutrition. *Cancer Epidemiology, Biomarkers and Prevention*, 19:2278-2286.
- Bursal, E. and Koksall, E.(2010) Evaluation of reducing power and radical scavenging activities of water and ethanol extracts from sumac (*Rhus coriaria* L.). *Food Research International*. Article in Press.
- Cai, Y. Z., & Corke, H. (1999). Amaranthus betacyanin pigments applied in model food systems. *Journal of Food Science*, 64,869–873.
- Cai, Y., Sun, M. and Corke, H. (2003). Antioxidant activity of Betalains from plants of the Amaranthaceae *Journal of Agricultural and Food Chemistry*., 51, 2288-2294.

- Cai, Y.Z., Sun, M. and Corke, H. (2005). Characterization and application of betalain pigments from plants of the Amaranthaceae. *Trends in Food Science & Technology*, 16 (9), 370-376. Pigments in Food, 3rd International Congress on Pigments in Food.
- Cao E.H., Xi-Qi Liu, Wang J-Jun Wang and Na-Fei Xu (1995). Evidence that lipid peroxidation products bind to DNA in liver cells. *Biochimica et Biophysica Acta* 1259, 187-191.
- Cao G, Sofic E, Prior RL (1996) Antioxidant capacity of tea and common vegetables. *Journal of Agricultural and Food Chemistry*, 44,3426–3430.
- Cao G.H, Sofic E. and Prior R.L. (1997). Antioxidant and prooxidant behaviour of flavanoids: Structure-activity relationships. *Free Radical Biology & Medicine*, Vol. 22(5), 749-760.
- Carr, A.C., Zhu, B.Z., Frei,B., (1999). Toward a new recommended dietary allowance for vitamin C based on antioxidant and health effects in humans. *American Journal of Clinical Nutrition*, 69(6), 1086-1107.
- Carr, A.C., Zhu, B.Z., Frei,B., (2000). Potential Antiatherogenic mechanism of ascorbate (Vitamin C) and α -tocopherol (Vitamin E). *Circulation Research*, 87, 349-354
- Castellar, R., Obon, J.M., Alacid, M. and Fernandez-Lopez, J.A.(2003). Color Properties and Stability of Betacyanins from *Opuntia* Fruits. *Journal of Agricultural and Food Chemistry*., 51,2772-2776.
- Cayuela, M.M (1994). Oxygen free radicals and human disease. *Biochimie*, 77, 147-161.
- Chanwitheesuk, A., Teerawutgulrag, A., & Rakariyatham, N. (2004). Screening of antioxidant compounds of some edible plants of Thailand. *Food Chemistry*, 3 (92), 491-497.
- Chee, Lee Lee (2003). Screening of Zingiberaceae extracts for antioxidant activities. *Tesis sarjana muda*. Universiti Malaya, Kuala Lumpur.
- Chen, Q-X. and Kubo, I. (2002). Kinetics of Mushroom Tyrosinase Inhibition by Quercetin. *Journal of Agricultural and Food Chemistry*., 50, 4108-4112.
- Chen, S.K., Tsai, M.L., Huang, J.R. and Chen, R.H.(2009). *In Vitro* Antioxidant Activities of Low-Molecular-Weight Polysaccharides with Various Functional Groups. *Journal of Agricultural and Food Chemistry*, 57, 2699-2704.
- Chiari, M.E., Joray, M.B., Ruiz, G., Palacios, S.M. and Carpinella, M.C. (2010). Tyrosinase inhibitory activity of native plants from central Argentina: Isolation of an active principle from *Lithrea molleoides*. *Food Chemistry* 120,10–14.
- Christian, G.D, (1994). *Analytical Chemistry* 5th Edition. John Wiley & Sons.

Christiana W., Barbara W., Katharina S., Harald S., Dietmar F., Ludwig B. (2005). *In vitro* Effects of Beet Root Juice on Stimulated and Unstimulated Peripheral Blood Mononuclear Cells. *American Journal of Biochemistry and Biotechnology* 1 (4): 180-185.

Collins, A.R. (2005). Antioxidant intervention as a route to cancer prevention. *European Journal of Cancer*, 41, 1923-1930.

Cook, N.C., Samman, S., (1996). Flavonoids-Chemistry, metabolism, cardioprotective effects, and dietary sources. *Journal of Nutritional Biochemistry*, 7, 69-76.

Czapski, J., Mikolajczyk, K. and Kaczmarek, M. (2009). Relationship between antioxidant capacity of Red Beet and contents of its betalain pigments. *Polish. Journal of Food and Nutritional Sciences.*, 59(2), 119-122.

Delgado-Vargas, F., Jimenez-Aparicio, A., & Paredes-Lopez, O. (2000). Natural pigments: Carotenoids, anthocyanins and betalains-Characteristics, biosynthesis, processing and stability. *CRC Critical Reviews in Food Science and Nutrition*, 40, 173-289.

Deng, J., Cheng, W. and Yang, G. (2011). A novel antioxidant activity index (AAU) for natural products using the DPPH assay. *Food Chemistry*, 125, 1430-1435.

Devalraju S., M.K. Arunasree, Karnati R., Reddy T.C., Gorla V. R., Pallu R. (2007). Betanin a betacyanin pigment purified from fruits of *Opuntia ficus-indica* induces apoptosis in human chronic myeloid leukemia Cell line-K562. *Phytomedicine*, 14, 739-746.

Dinan, L., Whiting, P. and Scott, A.J. (1998). Taxonomic distribution of phytoecdysteroids in seeds of members of the chenopodiaceae. *Biochemical Systematics and Ecology*, 26, 553-576.

Dmitrios, B. (2006). Sources of natural phenolic antioxidants. *Trends in Food Science & Technology*, 17, 505-512.

Duh, P-D., Yen, G-C., Yen, W-J., Wang, B-S. and Chang, L-W. (2004). Effects of Pu-erh Tea on Oxidative Damage and Nitric Oxide Scavenging. *Journal of Agricultural and Food Chemistry*, 52, 8169-8176.

Elisabeth (2007) Antioxidant properties of *Coriandrum sativum* and *Oenanthe javanica* leaves. *Tesis sarjana muda*. Universiti Malaya, Kuala Lumpur.

Geran, R.I., Greenberg, N.H., McDonald, M.M., Schumacher, A.M. and Abbott, B.J. (1972). Protocols for screening chemical agents and natural products against animal tumor and other biological systems. *Cancer Chemotherapy Reports*, 3, 17-19.

Schick, Yvonne Kerr, Proceeding in Hamilton Horizons student in College Seminar 235 (2008). Food for Thought: The Science, Culture, and Politics of Food.

- Epifano, F., Genovese, S., Menghini, L., Curini, M. (2007). Chemistry and pharmacology of oxyprenylated secondary plant metabolites. *Phytochemistry*, 68(7), 939-953.
- Escribano J., Pedreno, M.A., Garcia-Carmona, F. and Romualdo.(1998) Characterization of the antiradical activity of Betalains from *Beta vulgaris L.* roots. *Phytochemical Analysis*,9,124-127.
- Fong, S.Y. (2006). Antioxidant activities of *Hedyotis diffusa L.* and leaves of *Andrographis paniculata L.* Tesis sarjana muda. Universiti Malaya, Kuala Lumpur.
- Fotakis,G. and Timbrell, J.A. 2006). *In vitro* cytotoxicity assays: Comparison of LDH, neutral red, MTT and protein assay in hepatoma cell lines following exposure to cadmium chloride. *Toxicology Letters*,160,171–177.
- Frankel E.N. (1980).Lipid Oxidation.*Prog.Lipid.Pres*, 19, 1-22.
- Frankel E.N. (1999). Natural phenolic antioxidants and their impact on health. Antioxidant food supplements in human health. *Academic Press*, 25, 385-392.
- Freshney, R.I. (2000). Culture of animal cells: a manual of basic technique, 4th ed. Canada: Wiley-Liss.
- Fu, B., Li, H., Wang, X., Lee, F.S.C. and Cui, S. (2005). Isolation and Identification of Flavonoids in Licorice and a Study of Their Inhibitory Effects on Tyrosinase. *Journal of Agricultural and Food Chemistry*,53, 7408-7414.
- Gandia-Herrero, F., Escribano, J. and Garcia-Carmona, F. (2009). The role of Phenolic Hydroxy Groups in the Free Radical Scavenging Activity of Betalains. *Journal of Natural Products*, 72, 1142-1146.
- Gandia-Herrero, F., Escribano, J. and Garcia-Carmona, F. (2005). Betaxanthin as substrate for tyrosinase. An approach to the Role of Tyrosinase in the biosynthetic pathway of Betalains. *Plant Physiology*, 138 (1), 421-432.
- Georgiev, V., Ilieva, M., Bley, T. and Pavlow. (2008). Betalain production in plant in vitro systems. *Acta Physiologiae Plantarum*, 30.581–593.
- Georgiev, V.G.,Weber, J., Kneschke. E.M., Denev, P.N., Bley. T. and Pavlov, A.I. (2010). Antioxidant activity and phenolic content of Betalain extracts from intact plants and hairy root cultures of the red beetroot *Beta vulgaris* cv. Detroit Dark Red. *Plant Foods Hum Nutr*, 65(2):105-11.
- Geran, R.I., Greenberg, N.H., McDonald, M.M., Schumacher, A.M. & Abbott, B.J. (1972). Protocols for screening chemical agents and natural products against animal tumor and other biological systems. *Cancer Chemotherapy Reports*. 3(2): 17-19.
- Gil, M.I., Ferreres, F., Tomas-Barberan, F.A. (1998). Effect of modified atmosphere packaging on the flavonoids and vitamin C content of minimally processed Swiss chard

(*Beta vulgaris* subspecies *cycla*). *Journal of Agricultural and Food Chemistry*, 48, 2007-2012.

Gliszczynska-Swiglo, A., Szymusiak, H. and Malinowska, P. (2006). Betanin, the main pigment of red beet: Molecular origin of its exceptionally high free radical-scavenging activity. *Food Additives and Contaminants*, 23(11), 1079-1087.

Govinda, J.K., Magnus, A.A., Rajagopalan, S., Yoko, O., Atsushi, T., Eiichiro, I., Teruo, M., Midori, T., Takao, K., Hoyoku, N., Harukuni, T. (2003). Chemoprevention of DMBA-induced UV-B promoted, NOR-1-induced TPA promoted skin carcinogenesis, and DEN-induced phenobarbital promoted liver tumors in mice by extract of beetroot. *Pharmacological Research*, 47, 141-148.

Graham, R.D. and Ulrich, A. 1972. Potassium deficiency-induced changes in stomatal behavior, leaf water potentials, and root system permeability in *Beta vulgaris* L. *Plant Physiology*, 49, 105-109.

Gulcin, I. (2006). Antioxidant activity of caffeic acid (3,4-dihydroxycinnamic acid). *Toxicology*, 217, 213–220.

Gulcin, I. (2006). Antioxidant and antiradical activities of l-carnitine. *Life Sciences*, 78, 803 – 811.

Gulcin, I., Huyut, Z., Elmastas, M. and Aboul-Enein, H.Y. (2009). Radical scavenging and antioxidant activity of tannic acid. *Arabian Journal of Chemistry*, 3, 43-53.

Ha, T.J., Tamura, S. and Kubo, I. (2005). Effects of Mushroom Tyrosinase on Anisaldehyde. *Journal of Agricultural and Food Chemistry*, 53, 7024-7028.

Hakiman, M. and Maziah, M. (2009). Non enzymatic and enzymatic antioxidant activities in aqueous extract of different *Ficus deltoidea* accessions. *Journal of Medicinal Plants Research*, 3(3), 120-131.

Halliwell, B., Gutteridge, J.M.C., (1984). Oxygen toxicity, oxygen radicals, transitionmetals and disease. *Biochemistry Journal.*, 219, 1-14.

Halvorsen, B.L., Holte, K., Myhrstad, M.C.W. et al. (2002). A systematic screening of total antioxidants in dietary plants. *Journal of Nutrition*, 132, 461–471.

Hammerschmidt, P.A. and Pratt, D.E. 1978. Phenolic antioxidants of dried soybeans. *Journal of Food Science*. 43(2):556–559.

Han, X.H., Gao, Z.J. and Xiao, X.G. Enzymes and genes involved in the betalain biosynthesis in higher plants. *African Journal of Biotechnology*, 8(24), 6735-6744

Hanachi, P. and Golkho, S.H. (2009). Using HPLC to determination the composition and antioxidant activity of *Berberis vulgaris*. *European Journal of Scientific Research*, 1, 47-54.

Hani Fazlin Bt Mohd Fikry (2007). Antioxidant properties and essential oil components of *Piper sarmentosum*. *Tesis sarjana muda*. Universiti Malaya, Kuala Lumpur.

Harvey A.L. (2008). Review Natural products in drug discovery. *Drug Discovery Today*, 13,(19/20),894-901.

Harvey, A. Strategies for discovering drugs from previously unexplored natural products(2000). *Drug Discovery & Therapeutics*, 5(7), 294-300.

Hassimoto, N.M.A., Genovese, M.I. and Lajolo, F.M. (2005). Antioxidant activity of dietary fruits, vegetables and commercial frozen fruit pulps. *Journal of Agricultural and Food Chemistry*, 53, 2928-2935.

Hee Pyo, Y., Lee, T-C., Logendra, L. and Rosen, R.T. 2004. Antioxidant activity of phenolic compounds of Swiss chard (*Beta vulgaris* subspecies *cycla*) extracts. *Food Chemistry*, 85, 19-26.

Hong, W.K. and Sporn, M.B. (1997). Recent advances in chemoprevention of cancer: Frontiers in cancer research. *Science*, 278 (5340), 1073-1077.

Houghton, P., Lee, C.C. (2005). Cytotoxicity of plants from Malaysia and Thailand used traditionally to treat cancer. *Journal of Ethnopharmacology*, 100, 237-243.

Houghton, P.J., & Raman, A. 1998. *Laboratory handbook for the fractionation of natural extracts*, 1st Ed. London; Chapman and Hall.

Huang, D., Ou, B. and Prior, R.L. (2005). The chemistry behind antioxidant capacity assays. *Journal of Agricultural and Food Chemistry*, 53, 1841-1856.

Huey-Ko, H., Chang, W.L. and Lu, T.Z. (2008). Antityrosinase and antioxidant effects of *ent*-kaurane diterpenes from leaves of *Broussonetia papyrifera*. *Journal of Natural Products*, 71, 1930–1933.

Ikram, E.H.K., Eng, K.H., Jalil, A.M.M., Ismail, A., Idris, S., Azlan,A., Nazri, H.S.M., Diton, N.A.M., and Mokhtar, R.A.M. (2009). Antioxidant capacity and total phenolic content of Malaysian underutilized fruits. *Journal of Food Composition and Analysis*, 22, 388–393.

Ismail Dikici, Idris Mehmetoglu, Nebahat Dikici, Mehmet Bitirgen, Sevil Kurban (2005). Investigation of oxidative stress and some antioxidants in patients with acute and chronic viral hepatitis B and the effect of interferon- α treatment. *Clinical Biochemistry* 38, 1141-1144.

Ismail, A., Marjan, Z.M. and Foong, C.W. (2008). Total antioxidant activity and phenolic content in selected vegetables. *Food Chemistry*,87, 581–586.

- Ismail, H.I., Chan, K.W., Mariod, A.A. and Ismail, M. (2010). Phenolic content and antioxidant activity of cantaloupe (cucumis melo) methanolic extracts. *Food Chemistry*, 119, 643–647.
- Jastrebova, J., Wittho, C., Grahn, A., Svensson, U. and Gerstad, M.J. 2003. HPLC determination of folates in raw and processed beetroots. *Food Chemistry*, 80, 579–588.
- Javanmardi, J. Stushnoff, C., Locke, E. and Vivanco, J.M. (2003). Antioxidant activity and total phenolic content of Iranian Ocimum accessions. *Food Chemistry*, 83, 547–550.
- Jayakumar, T., Sakthivel, M., Thomas, P.A. and Geraldine, P. (2008). *Pleurotus ostreatus*, an oyster mushroom, decreases the oxidative stress induced by carbon tetrachloride in rat kidneys, heart and brain. *Chemico-Biological Interactions*, 176, 108–120.
- Jayaprakasha, G.T., Singh, R.P. and Sakariah, K.K. (2001). Antioxidant activity of grape seeds (*Vitis vinifera*) extracts on lipid peroxidation models in vitro. *Phytochemistry*, 73, 285–290.
- Jeong, S.H., Ryu, Y.B., Curtis-Long, M.J., Ryu, H.W., Baek, Y.S., Kang, J.E., Lee, W.S. and Park, K.H. (2009). Tyrosinase Inhibitory Polyphenols from Roots of *Morus Ihou*. *Journal of Agricultural and Food Chemistry*, 57, 1195–1203.
- Jerene K., (2007). Cytotoxic activity and anti-HPV of *Ganoderma tsugae* against selected cancer cell lines. *Unpublished Undergraduate Thesis*. University of Malaya, Kuala Lumpur.
- Jime´nez-Escrig, A., Dragsted, L. O., Daneshvar, B., Pulido, R., & Saura-Calixto, F. (2003). In vitro antioxidant activities of edible artichoke (*Cynara scolymus* L.) and effect on biomarkers of antioxidants in rats. *Journal of Agricultural and Food Chemistry*, 51, 5540–5545.
- Jiratanan, T. and Hai-Liu, R. (2004). Antioxidant Activity of Processed Table Beets (*Beta vulgaris* var. *conditiva*) and Green Beans (*Phaseolus vulgaris* L.). *Journal of Agricultural and Food Chemistry*, 52 (9), 2659–2670.
- Kahkonen, M.P., Hopia, A.I., Vuorela, H.J., Rauha, J.P., Pihlaja, K., Kujala, T.S., Heinonen, M. (1999). Antioxidant activity of plant extracts containing phenolic compounds. *Journal of Agricultural and Food Chemistry*, 47, 3954–3962.
- Kanner, J., Frankel, J.E., Granit, R., German, B., Kinsellatss, J.E. (1994). Natural antioxidants in grapes and wines. *Journal of Agricultural and Food Chemistry*, 42, 64–69.
- Kanner, J., Harel, S. and Granit, R. (2001). Betalains — A New Class of Dietary Cationized Antioxidants. *Journal of Agricultural and Food Chemistry*, 49 (11), 5178–5185.
- Kapadia GJ, Azuine MA, Sridhar R, Okuda Y, Tsuruta A, Ichiishi E, Mukainake T, Takasaki M, Konoshima T, Nishino H, Tokuda H. (2003). Chemoprevention of DMBA-

induced UV-B promoted, NOR-1-induced TPA promoted skin carcinogenesis, and DEN-induced phenobarbital promoted liver tumors in mice by extract of beetroot. *Pharmacological Research*, 47,141–148.

Kapadia GJ, Balasubramanian V, Tokuda H, Iwashima A, Nishino H (1997) Inhibition of 12-O-tetradecanoylphorbol-13-acetate induced Epstein-Barr virus early antigen activation by natural colorants. *Cancer Letters*, 115,173–178.

Kapadia, G.J., Tokuda, H., Konoshima, T., Nishino, H. (1996). Chemoprevention of lung and skin cancer by Beta vulgaris (beet) root extract. *Cancer Letters*, 100, 211-214.

Karou, D., Dicko, M.H., Simpore, J. and Traore, A.S. (2005). Antioxidant and antibacterial activities of polyphenols from ethnomedicinal plants of Burkina Faso. *African Journal of Biotechnology*, 4(8),823-828.

Karuppusamy, S. (2009). A review on trends in production of secondary metabolites from higher plants by *in vitro* tissue, organ and cell cultures. *Journal of Medicinal Plants Research*, 3(13), 1222-1239.

Keloff GJ.(2000). Perspectives on cancer chemoprevention research and drug development. *Advances in Cancer Research*,78,199-334.

Kim, H.J., Chen, F., Wang, X., Chung, H.Y. and Jin, Z. (2005). Evaluation of antioxidant activity of Vetiver (*Vetiveria zizanioides* L.) oil and identification of its antioxidant constituents. *Journal of Agricultural and Food Chemistry*, 53, 7691-7695.

King, R.J. (2000). *Cancer Biology*. Paperback Book. Prentice Hall Professional Technical Reference.

Koksal, E. and Gulcin, I. (2008). Antioxidant activity of cauliflower (*Brassica oleracea* L.). *Turkish Journal of Agriculture and Forestry*, 32, 65-78.

Krishnaiah, D., Sarbatly, R. and Bono, A. (2007). Phytochemical antioxidants for health and medicine – A move towards nature. *Biotechnology and Molecular Biology Review* ,1 (4),097-104.

Kubo, I. and Kinst-Hori, I. (1999,a). Flavonols from Saffron Flower: Tyrosinase Inhibitory Activity and Inhibition Mechanism. *Journal of Agricultural and Food Chemistry*, 47, 4121-4125.

Kubo, I. and Kinst-Hori, I.(1999,b). Tyrosinase inhibitory activity of the olive oil flavor compounds. *Journal of Agricultural and Food Chemistry*, 47, 4574-4578.

Kubola, J. and Siriamornpun, S. (2008). Phenolic contents and antioxidant activities of bitter melon (*Momordica charantia* L.) leaf, stem and fruit fraction extracts *in vitro*. *Food Chemistry*, 110, 881-890.

Kujala, T., Lojonen, J. and Pihlaja, K. (2001). Betalains and Phenolics in Red Beetroot (*Beta vulgaris*) Peel Extracts: Extraction and Characterisation. *Z. Naturforsch*, 56 (c), 343-348.

Kujala, T.S., Lojonen, J.M., Klika, K.D. and Pihlaja, K. (2000). Phenolics and Betacyanins in Red Beetroot (*Beta vulgaris*) Root: Distribution and effect of cold storage on the content of total phenolics and three individual compounds. *Journal of Agricultural and Food Chemistry*, 48, 5338-5342.

Kujala, T.S., Vienola, M.S., Klika, K.D., Lojonen, J.M., Pihlaja, K., 2002. Betalain and phenolic compositions of four beetroot (*Beta vulgaris*) cultivars. *European Food Research and Technology*, 214(6), 505-510.

Kulisic, T., Radonic, A., Katalinic, V. and Milos, M. (2004). Use of different methods for testing antioxidative activity of oregano essential oil. *Food Chemistry*, 85, 633-640.

Kuppusamy, U.R., Indran, & Balraj, B.R.S. (2002). Antioxidant effects of local fruits and vegetable extracts. *Journal of Tropical Medicinal Plants*, 3(1), 47-53.

Laguerre M., Lecomte J., Villeneuve P. (2007). Evaluation of the ability of antioxidants to counteract lipid oxidation: Existing methods, new trends and challenges. *Progress in Lipid Research* 46, 244-282.

Lange, W., Brandenburg, W.A. and De Bock, T.S.M. (1999). Taxonomy and cultonony of Beet (*Beta vulgaris* L.). *Botanical Journal of the Linnean Society*, 130, 81-96.

Lee, B.J., Kim, Y. and Cho, M.H. (1999). Effects of phytic acid content, storage time and temperature on lipid peroxidation in muscle foods. *Journal of Food Hygiene and Safety*, 14 (1), 27-33.

Lee, H.W. (2006). Antioxidant activities of leaves of *Ficus religiosa* and *Cardiospermum halicacabum*. *Tesis sarjana muda*. Universiti Malaya, Kuala Lumpur.

Leland, J.C, Kirakosyan, A., Kaufman, P.B., Warber, S.L., Duke, J.A., Briemann, H.L. (2006). *Natural Products from Plants*, Second Edition. Taylor and Francis Group.

Letelier, M.E., Berridos, A.M., Cortes-Tronsoco, J., Jara-Sandoval, J., Holst, M., Palma, K, Montoya, M., Miranda, D. and Gonzalez-Lira, V.(2008). DPPH and oxygen free radicals as pro-oxidant of biomolecules. *Toxicology in Vitro*, 22, 279-286.

Lim, Y.Y. and Quah, E.P.L. (2007). Antioxidant properties of different cultivars of *Portulaca oleraceae*. *Food Chemistry*, 103, 734-740.

Ling Jackie Chooi Lin. (2006). *Chemical and antioxidant investigation on essential oil of Zingiber officinale var. rubrum* (jahe emprit). *Tesis sarjana muda*. Universiti Malaya, Kuala Lumpur.

- Ling, L.T., Radhakrishnan, A.K., Subramaniam, T., Cheng, H.M. and Palanisamy, U.D. (2010). Assessment of Antioxidant Capacity and Cytotoxicity of Selected Malaysian Plants. *Molecules*, 15, 2139-2151.
- Linscheid, M. and Westmoreland, D.G. (1994). Application of liquid chromatography – mass spectrometry. *Pure & Applied Chemistry*, 66(9), 1913-1930.
- Liu X., Ardo, S., Bunning, M., Parry, J., Zhou, K., Stushnoff, C., Stoniker, F., Yu, L. and Kendall, P. (2007). Total phenolic content and DPPH radical scavenging activity of lettuce (*Lactuca sativa* L.) grown in Colorado. *Lebensmittel-Wissenschaft und-Technologie*, 40, 552–557.
- Liu, Q., Zhang, Y-J., Yang, C-R. and Xu, M. (2009). Phenolic antioxidants from green tea produced from *Camellia crassicolumna* var. *multiplex*. *Journal of Agricultural and Food Chemistry*, 57, 586-590.
- Lu, F., & Foo, L. Y. (1995). Phenolic antioxidant component of evening primrose. In A. S. H. Ong, E. Niki, & L. Packer (Eds.), *Nutrition, lipids, health and disease*. Campaign: American Oil Chemists Society Press.
- Lue, B-M., Nielsen, N.S., Jacobsen, C., Hellgren, L., Xu, Z.G.X. (2010). Antioxidant properties of modified rutin esters by DPPH, reducing power, iron chelation and human low density lipoprotein assays. *Food Chemistry*, 123, 221-230.
- Mabry, T. J., & Dreiding, A. S. (1968). The betalains. In T. J. Mabry, R. E. Alston, & V. C. Runeckles (Eds.), *Recent advances in phytochemistry*, 1, 145-160.
- Magalhaes, L.M., Segundo, M.A., Reis, S., Lima, J. (2008). Methodological aspects about *in vitro* evaluation of antioxidant properties. *Analytica Chimica Acta*, 613(1), 1-19.
- Maisuthisakul, P. and Gordon, M.H. (2009). Antioxidant and tyrosinase inhibitory activity of mango seed kernel by product. *Food Chemistry*, 117, 332-341.
- Maisuthisakul, P., Suttajit, M. and Pongsawatmanit, R. (2007). Assessment of phenolic content and free radical-scavenging capacity of some Thai indigenous plants. *Food Chemistry*, 100, 1409-1418.
- Maizatul. (2008). Cytotoxic and human papiloma virus activities in selected vegetables. *Unpublished Masters Thesis*. Universiti Malaya, Kuala Lumpur.
- Mansukhlal, C.W., Taylor, H.L., Wall, M.E., Coggon, P. and McPhail, A.T. (1971). Plant antitumor agents. VI. Isolation and structure of taxol, a novel antileukemic and antitumor agent from *Taxus brevifolia*. *Journal of the American Chemical Society*, 93(9), 2325-2327.
- Martin, S.S. (1977). Accumulation of the flavonoids betagarin and betavulgarin in *Beta vulgaris* infected by the fungus *Cercospora beticola*. *Physiological Plant Pathology*, 11(3), 297-303.

- Martin, S.S. (1989). Analysis of Constitutive and Induced Phenolics of *Beta vulgaris* by High Performance Liquid Chromatography. *Analysis of Constitutive and Induced Phenolics*, 26(2), 33-39.
- Masamoto, Y., Ando, H., Murata, Y., Shimoishi, Y., Tada, M. and Takahata, K. (2003). Mushroom Tyrosinase Inhibitory Activity of Esculetin Isolated from Seeds of *Euphorbia lathyris* L. *Bioscience, Biotechnology and Biochemistry*, 67(3),631-634.
- Masuda, T., Yamashita, D., takeda, Y., and Yonemori, S. (2005). Screening for tyrosinase inhibitors among extracts of seashore plants and identification of potent inhibitors from *Garcinia subelliptica*. *Bioscience, Biotechnology and Biochemistry*, 69(1),197-201.
- Mates, J.M., Gomez, C.P., Castro, I.N. (1999). Antioxidants enzymes and human diseases. *Clinical Biochemistry*, 32(8), 595-603.
- Matsuura, R., Ukeda, H. and Sawamura, M. (2006). Tyrosinase inhibitory activity of citrus essential oils. *Journal of Agricultural and Food Chemistry*, 54, 2309-2313 .
- McChesney, J.D., Venkataraman, S.K., & Henri, J.T. (2007). Plant natural products: Back to the future or into extinction? *Phytochemistry*, 68, 2015-2022.
- McCord, J.M. and Fridovich, I. (1969). Superoxide Dismutase, An enzymic function for erythrocyte (Hemocuprein). *The Journal of Biological Chemistry*, 244 (22), 6049-6055.
- Mee, S.P.H., Lee, V., Baldwin, J.E. and Cowley,A. 2004. Total synthesis of 5,5',6,6'-tetrahydroxy-3,30-biindolyl, the proposed structure of a potent antioxidant found in beetroot (*Beta vulgaris*). *Tetrahedron*, 60, 3695-3712.
- Miraliakbari, H. and Shahidi, F. (2008). Antioxidant activity of minor components of tree nut oils. *Food Chemistry*, 111, 421–427.
- Miyazawa, M., Oshima, T., Koshio, K., Itsuzaki, Y. and Anzai, J. (2003). Tyrosinase Inhibitor from Black Rice Bran. *Journal of Agricultural Food Chemistry*, 51, 6953-6956.
- Molyneux, P., (2004). The use of the stable free radical diphenylpicryl-hydrazyl (DPPH) for estimating antioxidant activity. *Songklanakarin Journal of Science and Technology*, 26(2), 211-219.
- Moreno, C.S. (2002). Review: Methods Used to Evaluate the Free Radical Scavenging Activity in Foods and Biological Systems. *Food Science and Technology International*, 8, 121-137.
- Muhammad Muzhafar Mohamad Noor (2007). Antioxidant effect of crude extract of daun kadok (*Piper sarmentosum*). *Tesis sarjana muda*. Universiti Malaya, Kuala Lumpur.

- Naidu, J. (2008). Antioxidant activity of *Melicope ptelefolia* (tenggek burung). *Tesis sarjana muda*. Universiti Malaya, Kuala Lumpur.
- Nascimento, F.R.F., Cruz, G.V.B., Pereira, P.V.S., Maciel, M.C.G., Silva, L.A., Azevedo, A.P.S., Barroqueiro, E.S.B., Guerra, R.N.M. (2006). Ascitic and solid Ehrlich tumor inhibition by *Chenopodium ambrosioides* L. treatment. *Life Sciences*, 78, 2650-2653.
- Nascimento, M.A., Silva, A.K., Franc, L.C.B., Quignard, E.L.J., Lopez, J.A. and Almeida, M.G. (2006). *Turnera ulmifolia* L. (Turneraceae): Preliminary study of its antioxidant activity. *Bioresource Technology*, 97, 1387–1391.
- Netzel M., Stintzing F.C., Quaaas D., Strass G., Carle R., Bitsch I., Frank T. (2005) Renal excretion of antioxidative constituents from red beet in humans. *Food Research International*, 38, 1051—1058.
- Ng L.T., Ko H. and Lu T.M. (2009). Potential antioxidants and tyrosinase inhibitors from synthetic polyphenolic deoxybenzoins. *Bioorganic & Medicinal Chemistry*, 17, 4360-4366.
- Niki, E. (2010). Assessment of Antioxidant Capacity in vitro and in vivo. *Free Radical Biology & Medicine*, 49 ,503–515.
- Niki, E., Yoshida, Y., Saito, Y. and Noguchi, N. (2005). Lipid peroxidation: Mechanisms, inhibition, and biological effects. *Biochemical and Biophysical Research Communications*, 338, 668–676.
- Ninfali, P., Bacchiocca, M., Antonelli, A., Biagiotti, E., Di Gioacchino, A.M., Piccoli, G., Stocchi, V., Brandi, G. (2007). Characterization and biological activity of the main flavonoids from Swiss Chard (*Beta vulgaris* subspecies *cycla*). *Phytomedicine*, 14, 216-221.
- Oliveira, T.D., Nagem, T.J. Costa, R.D.M., Marciano, D.C.L., Magalhaed, N.M., Stringheta, P.C, Queiroga D.L.E., Kling, D.M.G.H and Silva, V.H. (2004). Biological properties of natural dyes. *Ars Pharmaceutica*, 45(1), 5-20.
- Osawa,T., Namiki, M., (1980). A novel type of antioxidant isolated from leaf wax of Eucalyptus leaves. *Agricultural and Biological Chemistry*, 45(3), 735-739.
- Ou, B., Huang, D., Hampsch-Woodill, M., Flanagan, J.A. & Deemer, E.K. (2002). Analysis of antioxidant activities of common vegetables employing oxygen radical absorbance capacity (ORAC) and ferric reducing antioxidant power (FRAP) assays: a comparative study. *Journal of Agricultural and Food Chemistry*, 50, 3122–3128.
- Oyaizu, M. (1986).Studies on product of browning reaction prepared from glucose amine. *Japanese Journal of Nutrition*, 44, 307–315.

- Parejo, I., Codina, C., Petrakis, C. and Kefalas, P. (2000). Evaluation of scavenging activity assessed by Co(II)/EDTA-induced luminal chemiluminescence and DPPH free radical assay. *Journal of Pharmacological and Toxicological Methods*, 44, 507-512.
- Pavlov A, Georgiev V, Ilieva M (2005) Betalain biosynthesis by red beet (*Beta vulgaris* L.) hairy root culture. *Process Biochemistry* 40,1531–1533.
- Pavlov A, Georgiev V, Kovatcheva P (2003) Relationship between type and age of the inoculum cultures and betalains biosynthesis by *Beta vulgaris* hairy root culture. *Biotechnol Letters*, 25,307–309.
- Pavlov A, Kovatcheva P, Georgiev V, Koleva I, Ilieva M (2002) Biosynthesis and radical scavenging activity of betalains during the cultivation of red beet (*Beta vulgaris*) hairy root cultures. *Z Naturforsch C*, 57,640–644.
- Pedreno, M.A. and Escribano, J. (2000). Studying the oxidation and the antiradical activity of betalain from beetroot. *Journal of Biological Education*, 35(1), 49-51.
- Piatelli, M. (1981). The betalains: Structure, biosynthesis, and chemical taxonomy. *The Biochemistry of Plants*, 7, 557-575.
- Pitalua, E., Jimenez, M., Vernon-Carter, E.J. and Beristain, C.I. 2010. Antioxidative activity of microcapsules with beetroot juice using gum Arabic as wall material. *Food and bioproducts processing*, 88, 253-258.
- Prakash, A., Rigelhof, F., Miller, E. (2001). Antioxidant activity. *Medillion Laboratories Analytical Progress*, 19(2).
- Prior, R.L. and Cao, G.(2000). Analysis of Botanicals and Dietary Supplements for Antioxidant Capacity: A Review. *Journal of AOA international*, 83(4), 950-956.
- Prior, R.L., Wu, X. and Schaich, K. (2005). Standardized Methods for the Determination of Antioxidant Capacity and Phenolics in Foods and Dietary Supplements. *Journal of Agricultural and Food Chemistry*, 53, 4290-4302.
- Rael, T.L., Thomas, W.G., Craun, M.L., Curtis, G.C., Bar-Or, R., Bar-Or, D., (2004). Lipid peroxidation and thiobarbituric acid assay: Standardization of the assay when using saturated and unsaturated fatty acids. *Journal of Biochemistry and Molecular Biology*, 37(6), 749-752.
- Ramanathan, N., Osawa, T., Hiroto, O. and Kawakishi, S. (1995). The contribution of plant food antioxidants to human health. *Trends in Food Science & Technology*, 6, 75-82.
- Reddy L., B. Odhav, K.D. Bhoola. (2003). Natural products for cancer prevention: a global perspective. *Pharmacology and Therapeutics*, 99,1– 13.

- Reddy, M.k., Alexander-Lindo, R.I., Nair, M.G., (2005). Relative inhibition of lipid peroxidation, cyclooxygenase enzymes, human tumor cell proliferation by natural food clors. *Journal of Agricultural and Food Chemistry*, 53, 9268-9273.
- Reddy,L., Odhav, B. and Bhoola, K.D. (2003). Natural products for cancer prevention: a global perspective. *Pharmacology & Therapeutics* ,99,1– 13.
- Rey, A.I., Hopia, A., Kivikari, R. and Kahkonen, M. (2005). Use of natural food/plant extracts: cloudberry (*Rubus Chamaemorus*), beetroot (*Beta Vulgaris* “*Vulgaris*”) or willow herb (*Epilobium angustifolium*) to reduce lipid oxidation of cooked pork patties. *Lebensmittel-Wissenschaft und-Technologie*, 38, 363-370.
- Reynoso, R., Garcia, F.A., Morales, D. and Gonzalez de Mejia, E.(1997). Stability of Betalain Pigments from a Cactacea Fruit. *Journal of Agricultural and Food Chemistry*, 45, 2884-2889.
- Rufino, M.D.S.M., Alves,R.E., de Brito, E.S., Jimenez, J.P., Calixto, F.S. Filho, J.M. (2010). Bioactive compounds and antioxidant capacities of 18 non-traditional tropical fruits from Brazil. *Food Chemistry* ,121, 996–1002.
- Rufino, M.S.M., Fernandez, F.A.N., Alves, R.E. and de Brito, E. S. (2009). Free radical-scavenging behaviour of some north-east Brazilian fruits in a DPPH system. *Food Chemistry*, 114, 693–695.
- Sawangjaroen, N., Sawangjaroen,K., Poonpanang, P., (2004). Effects of *Piper longum* fruit, *Piper sarmentosum* root and *Quercus infectoria* nut gall on caecal amoebiasis in mice. *Journal of Ethnopharmacology*, 91, 357-360.
- Sen, S., Chakraborty, R., Sridhari, C., Reddy, Y.S.R. and De, B. (2010). Free radicals, antioxidants, diseases and phytomedicines: Current status and future prospect. *International Journal of Pharmaceutical Sciences Review and Research*, 3(1), 91-100.
- Sepu'lveda-Jime'nez G, Rueda-Ben'i'tez P, Porta H, Rocha-Sosa M (2004) Betacyanin synthesis in red beet (*Beta vulgaris*) leaves induced by wounding and bacterial infiltration is preceded by an oxidative burst. *Physiological and Molecular Plant Pathology*, 64,125–133.
- Sharififar, F., Nudeh, G.D. and Mirtajaldini, M. (2009). Major flavonoids with antioxidant activity from *Teucrium polium* L. *Food Chemistry* ,112, 885–888.
- Sharma, O.P. and Bhat, T.K. DPPH antioxidant assay revisited (2009). *Food Chemistry* 113, 1202–1205.
- Sheu, J.Y., Chen, C.Y., Tsai, L.Y., and Huang, Y.L. (2003). Spectrometric determination of a thiobarbituric acid – reactive substance in human hair. *Analytical Sciences*, 19, 957-960.

- Shimada, K., Fujikawa, K., Yahara, K., & Nakamura, T. (1992). Antioxidative properties of xanthan on the autoxidation of soybean oil in cyclodextrin emulsion. *Journal of Agricultural and Food Chemistry*, 40, 945–948.
- Shukla, S., Mehta, A., Bajpai, V.K. and Shukla, S. (2009). In vitro antioxidant activity and total phenolic content of ethanolic leaf extract of *Stevia rebaudiana* Bert. *Food and Chemical Toxicology*, 47, 2338–2343.
- Siddhuraju, P., (2005) The antioxidant activity and free radical-scavenging capacity of phenolics of raw and dry heated moth bean (*Vigna aconitifolia*) (Jacq.) Marechal seed extracts. *Food Chemistry*, 99 (2006), 149–157.
- Sim, K.S. (2010). Bioactivity and chemical investigations of *Pereskia bleo* and *Pereskia grandilifolia*. *Tesis doktor falsafah*. Universiti Malaya, Kuala Lumpur.
- Simon P. H. Mee, Victor Lee, Jack E. Baldwin, and Andrew Cowley. (2004). Total synthesis of 5',5'',6',6''-tetrahydroxy-3,30-biindolyl, the proposed structure of a potent antioxidant found in beetroot (*Beta vulgaris*). *Tetrahedron*, 60, 3695–3712.
- Singleton, V. L., & Rossi, J. A. (1965). Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *American Journal of Enology and Viticulture*, 16, 144–158.
- Son, S. and Lewis, B.A. Free radical scavenging and antioxidative activity of caffeic acid amide and ester analogues: structure-activity relationship.(2002). *Journal of Agricultural and Food Chemistry*, 50 (3), 468-472.
- Sreekanth D, Arunasree MK, Roy KR, Chandramohan T, Reddy GV, Reddanna P (2007) Betanin a betacyanin pigment purified from fruits of *Opuntia ficus-indica* induces apoptosis in human chronic myeloid leukemia Cell line-K562. *Phytomedicine*, 14, 739–746.
- Srivastava, A., Harish, S.R. and Shivanandappa, T. (2006). Antioxidant activity of the roots of *Decalepis hamiltonii* (Wight & Arn.). *Lebensmittel-Wissenschaft und-Technologie*, 39, 1059–1065.
- Stangeland, T., Wangensteen, H., Katuura, E., Lye, K.A. and Paulsen, B.S. (2010). Antioxidant and anti-plasmodial activity of extracts from three Ugandan medicinal plants. *Journal of Medicinal Plants Research*, 4 (18), 1916-1923.
- Steiner, U., Schliemann, W., Boehm, H. and Strack, D. (1999). Tyrosinase involved in betalain biosynthesis of higher plants. *Planta*, 208, 114-124.
- Stephen, N. (2004). Beetroot E- Book.
- Stintzing FC, Carle R (2004) Functional properties of anthocyanins and betalains in plants, food, and in human nutrition. *Trends in Food Science and Technology*, 15, 19–38.

- Stintzing FC, Carle R (2007) Betalains—emerging prospects for food scientists. *Trends in Food Science and Technology*, 18,514–525.
- Stintzing, C.F., Conrad, J., Klaiber, I., Beifuss, U., Carle, R. (2004). Structural Investigations on betacyanin pigments by LC NMR and 2D NMR spectroscopy. *Phytochemistry* 65,415-422.
- Stintzing, F.C. and Carle, R.(2004). Functional properties of anthocyanins and betalains in plants, food, and in human nutrition. *Trends in Food Science & Technology* 15, 19–38.
- Stintzing, F.C., Schieber, A. and Carle, R.2002. Identification of betalains from Yellow Beet (*Beta vulgaris* L.) and Cactus Pear [*Opuntia ficus-indica* (L)Mill] by High-Performance Liquid Chromatography-Electrospray Ionization Mass Spectrometry. *Journal of Agricultural and Food Chemistry*, 50, 2302-2307.
- Strack D, Vogt T, Schliemann W (2003) Recent advances in betalain research. *Phytochemistry* 62:247–269.
- Strack, D., Engel, U. and Wray, V. (1987). Neobetanin: A new plant constituent. *Phytochemistry*, 26(8), 2399-2400.
- Steele, V.E. and Kelloff, G.J. (2005). Development of cancer chemopreventive drugs based on mechanistic approaches. *Mutation Research*, 591, 16-23.
- Stryer, L., Tymoczko, J.L., Berg, J.M., (2003). *Biochemistry*, 5th edition. W.H. Freeman and Company, New York.
- Suja, K.P., Jayalekshmy, A. and Arumughan, C. (2005). Antioxidant activity of sesame cake extract. *Food Chemistry* ,91, 213–219.
- Sun, T. and Ho, C.H. (2005). Antioxidant activities of buckwheat extracts. *Food Chemistry*,90, 743–749.
- Sun, T., Powers, J.R., and Tang, J. (2007). Evaluation of the antioxidant activity of asparagus,broccoli and their juices. *Food Chemistry*,105,101–106.
- Tachakittirungrod, S., Okonogi, S., Chowwanapoonpohn, S. (2006). Study on antioxidant activity of certain plants in Thailand: Mechanism of antioxidant action of guava leaf extract. *Journal of Food Chemistry*, 103 (2), 381-388.
- Taga, M. S., Miller, E. E., Pratt, D. E. (1984).Abstract. Chia seeds as a source of natural lipid antioxidants. *Journal of American Oil Chemist's Society*, 61, 928-931.
- Tan, G.W. (2006). Antioxidant activity of *Oxalis corniculata* and leaves of *Cucurbita moschata*. *Tesis sarjana muda*. Universiti Malaya, Kuala Lumpur.

- Tawaha, K., Alali, F.Q., Gharaibeh, M., Mohammad, M., El-Elimat, T. (2007). Antioxidant activity and total phenolic content of selected Jordanian plant species. *Food Chemistry*, 104, 1372–1378.
- Tenpe, C.R., Upaganlawar, A., Bhagat A., Yeole, P.G., 2008. *In-vitro* antioxidant and free radical scavenging activity of *Jasminum sambac* Linn. leaves. *Pharmacognosy Magazine*, 4(15(suppl)), 124-129.
- Thaipong, K., Boonprakob, U., Crosby, K., Cisneros-Zevallos, L. and Byrne, D.H. (2006). Comparison of ABTS, DPPH, FRAP, and ORAC assays for estimating antioxidant activity from guava fruit extracts. *Journal of Food Composition and Analysis*, 19, 669-675.
- Tiwari, A.K. (2008). Imbalance in antioxidant defence and human diseases: Multiple approach of natural antioxidants therapy. *CURRENT SCIENCE*, 9, 81, 1179-1187.
- Towatana, H.N., Chaiyamutti, P., Panthong, K., Mahabusarakam, W., Rukachaisirikul, V. (2006). Antioxidative and free radical scavenging activities of some plants used in Thai folk medicine. *Journal of Pharmaceutical Biology*, 44 (3), 221-228.
- Trejo-Tapia, G., Balcazar-Aguilar, J.B., Martinez-Bonfil, B., Salcedo-Morales, G., Jaramillo-Flores, M., Arenas-Ocampo, M.L., Jimenez-Aparicio, A. (2008). Effect of screening and subculture on the production of betaxanthins in *Beta vulgaris* L. var. 'Dark Detroit' callus culture. *Innovative Food Science & Emerging Technologies*, 9, 32-36.
- Tulp, M., Bohlin, L. (2004). Unconventional natural sources for future drug discovery. *Drug Discovery and Therapeutics*, 10 (9), 450-458.
- Ukrisdawithid, S., Sukmomtip, M., Bhusawang, P. and Pongsamart, S. (2008). A topical gel contained extract of seed coat of *Tamarindus indica* and its *in-vitro* antilipid peroxidation. Proceedings of 34th Congress on Science and Technology of Thailand, 1-6.
- Valko, M., Leibfritz, D., Moncol, J., Cronin, M.T.D., Mazur, M., Telser, J. (2006). Free radicals and antioxidants in normal physiological functions and human disease. *Journal of Biochemistry & Cell Biology*, 39(1), 44-84.
- Valvi, S.R., Rathod, V.S., and Yesane, D.P. (2011). Screening of three wild edible fruits for their antioxidant potential. *Current botany*, 2(1), 48-52.
- Velioglu, Y.S., Mazza, G., Gao, L., Oomah, B.D. (1998). Antioxidant activity and total phenolics in selected fruits, vegetables and grain products. *Journal of Agricultural and Food Chemistry*, 46, 4113-4117.
- Vimala, S., Mohd. Ilham Adenan, Abdull Rashih Ahmad & Sahdan, R. (2003). Natural antioxidants: *Piper sarmentosum* (Kadok) and *Morinda elliptica* (Mengkudu). *Tesis Doktor Falsafah*. Universiti Malaya, Kuala Lumpur.

- Vimala, S., Mohd. Ilham Adenan, Abdull Rashih Ahmad & Sahdan, R. (2003). Natural antioxidants : *Piper sarmentosum* (Kadok) and *Morinda elliptica* (Mengkudu). *Malaysian Journal Nutritional*, 9 (1), 41-45.
- Vimala, S., Norhanom, A.W. and Yadav, M. (1999). Anti-tumour promoter activity in Malaysian ginger rhizobia used in traditional medicine. *British Journal of Cancer*, 80 (1/2), 110-116.
- Vinson, J.A., Dabbagh, Y.A., Serry, M.M., Jang, J.(1995). Plant flavonoids, especially tea flavonols, are powerful antioxidants using an *in vitro* oxidation model for heart disease. *Journal of Agricultural and Food Chemistry*, 43, 2800-2803.
- Vinson, J.A., Hao, Y., Su, X. & Zubik, L. (1998). Phenol antioxidant quantity and quality in foods: vegetables. *Journal of Agricultural and Food Chemistry*, 46, 3630–3634.
- von Elbe JH, Schwartz SJ (1981) Absence of mutagenic activity and short-term toxicity study of beet pigments as food colorants. *Archives of Toxicology* 49:93–98.
- von Elbe, J.H. (1975). Stability of Betalaines as food colors. *Food Technology*, 42-46.
- Wang, T., Jonsdottir, R., Olafsdottir, G. (2009). Total phenolic compounds, radical scavenging and metal chelation of extracts from Icelandic seaweeds. *Food Chemistry*, 116, 240-248.
- Wall ME,Wani MC, Cook CE,Palmer KH, McPhail AT, Sim GA. (1966) Plant anti-tumor agents. The isolation and structure of camptothecin, a novel alkaloidal leukemia and tumor inhibitor from *Camptotheca acuminata*. *Journal of American Chemical Society*,88,3888-3890.
- Wettasinghe M, Bolling B, Plhak L, Xiao H, Parkin K (2002b) Phase II enzyme-inducing and antioxidant activities of beetroot (*Beta vulgaris L.*) extracts from phenotypes of different pigmentation. *Journal of Agricultural and Food Chemistry*, 50, 6704–6709.
- Wettasinghe, M., and Shahidi, F. (2000). Scavenging of reactive-oxygen species and DPPH free radicals by extracts of borage and evening primrose meals. *Food Chemistry*, 70,17-26.
- Weyermann, J.,Lochmann, D., Zimmer, A.(2005). A practical note on the use of cytotoxicity assays. *International Journal of Pharmaceutics*,288,369–376.
- Wilson, A.P. (1986). Cytotoxicity and viability assays in animal cell culture – a practical approach. Edited by Freshney, R.I. IRL Press Limited, New York
- Winkler, C., Wirleitner, B., Schroecksnadel, K., Schennach, H. and Fuchs, D. (2005). *In vitro* Effects of Beet Root Juice on Stimulated and Unstimulated Peripheral Blood Mononuclear Cells. *American Journal of Biochemistry and Biotechnology*, 1(4), 180-185.

Xuan Liu , Yanxiang Gao , Xiaoting Peng , Bin Yang , Honggao Xu , Jian Zhao (2008). Inactivation of peroxidase and polyphenol oxidase in red beet (*Beta vulgaris* L.) extract with high pressure carbon dioxide. *Innovative Food Science Emerging Technologies*, 9(1)-24-33.

Young, H.P., Tung, C.L., Logendra, L., Rosen, R.T. (2004). Antioxidant activity and phenolic compounds of Swiss chard (*Beta vulgaris* subspecies *cycla*) extracts. *Food Chemistry*, 85, 19-26.

Yuan, Y.V., Bone, D.E. and Carrington, M.F. (2005). Antioxidant activity of dulse (*Palmaria palmata*) extract evaluated in vitro. *Food Chemistry*, 91, 485–494.

Zakharova, N. S.; Petrova, T. A.(1998) Relationship between the structure and antioxidant activity of various betalains. *Prikl.Biochim. Mikrobiol.*, 34, 199-202.

Zhang, Dong-Lin, & Hamauzu, Y. (2003). Phenolic compounds, ascorbic acid, carotenoids and antioxidant properties of green, red and yellow bell peppers. *Journal of Food, Agriculture and Environment*, 1, 22–27.

Zhong Cai, Y., Sun, M. and Corke, H. Characterization and application of betalain pigments from plants of the Amaranthaceae. *Trends in Food Science & Technology*, 16(9), 370-376.

APPENDIX 1: LIST OF PAPERS PRESENTED

1. Poster presentation at 2nd International Symposium on Medicinal and Nutraceutical Plants (2nd ISMNP)
Title: Antioxidant Activity of *Beta vulgaris* (Beetroot)
Venue: All Indian Institute of Medical Sciences (AIIMS), New Delhi, India
Date: 25-27 November 2009
2. Poster presentation at International Conference on Natural Products (ICNP)
Title: Comprehensive Study on the Antioxidant Property of *Beta vulgaris* (Beetroot)
Venue: Bayview Beach Resort, Batu Ferringhi, Penang
Date: 11-12 December 2010
3. Oral presentation at 15th Biological Sciences Graduate Congress (BSGC)
Title: Detailed Study on the Antioxidant Property of *Beta vulgaris* (Beetroot)
Venue: Institute of Biological Sciences, University of Malaya, Kuala Lumpur
Date: 15-17 December 2010
4. Poster presentation at One day scientific and motivation seminar by Cenar (Centre for natural products and drug discovery, UM)
Title: Comprehensive Study on the Antioxidant Property of *Beta vulgaris* (Beetroot)
Venue: Crystal Crown Hotel, Petaling Jaya, Kuala Lumpur
Date: 22nd December 2010
5. Poster presentation at 25th Scientific Meeting of Malaysian Society of Pharmacology and Physiology (25th MSPP)
Title: Evaluation of Lipid Oxidation Inhibition Activity of *Beta vulgaris* L. (beetroot)
Venue: University Putra Malaysia (UPM)
Date: 25-26 May 2011
6. Poster presentation at 16th Biological Sciences Graduate Congress (16th BSGC)
Title: Evaluation of Lipid Oxidation Inhibition Activity of *Beta vulgaris* L. (beetroot).
Venue: National University of Singapore (NUS)
Date: 12-14th Dec 2011

APPENDIX 2: Preparation of Reagents

1. Antioxidant Assays

DPPH (1,2-dipheyl-2-picrylhydrzyl) Radical Scavenging Assay

Ascorbic Acid / Butylated Hydroxyanisole (BHA)

A stock of ascorbic acid/BHA (Sigma) in methanol was prepared at a concentration of 400 µg/ml by diluting 0.04 g ascorbic acid/BHA in 100 ml of methanol. The stock solution was kept in Duran bottle wrapped with aluminium foil. This is to ensure that there is no degradation activity as it is easily degraded in the presence of light (light sensitive compound). The preparation of the stock is also conducted in a dark place.

1,2-Diphenyl-2-picrylhydrzyl (DPPH)

A stock of DPPH (Sigma) in methanol was prepared at concentration of 8mg/ml (0.08g + 10 ml of methanol). The stock solution was kept in Duran bottle wrapped with aluminium foil. This is to ensure that there is no degradation activity as it is easily degraded in the presence of light (light sensitive). The preparation of stock is also conducted in a dark room.

2. Reducing Power Assay

Potassium Ferricyanide (1%,w/v) $K_3Fe(CN)_6$

Potassium Ferricyanide 1% was prepared by dissolving 0.1 gm of Potassium Ferricyanide (Sigma) in 10ml distilled water. The solution was kept in centrifuge tube and wrapped with aluminium foil.

Trichloroacetic acid (10%, w/v)

TCA 10% was prepared by dissolving 5gm of TCA (Sigma) in 500ml distilled water. The solution was kept in flask wrapped with aluminium foil.

Ferric Chloride (0.1%,w/v) $FeCl_3.6H_2O$

Ferric chloride solution was prepared by dissolving 0.01gm of ferric chloride (Sigma) in 10ml distilled water. The solution was kept in centrifuge tube and wrapped in aluminium foil.

0.2M Phosphate Buffer pH6.6

i. 0.2M Monobasic Stock

0.2M Monobasic Stock was prepared by dissolving 27.80gm of sodium phosphate monobasic (NaH_2PO_4 , Sigma) in 1000ml distilled water.

ii. 0.2M Dibasic Stock

0.2M Dibasic Stock was prepared by dissolving 53.65gm of sodium phosphate monobasic (Na_2HPO_4 , Sigma) in 1000ml distilled water.

iii. 0.2M Phosphate Buffer pH6.6

0.2M Phosphate Buffer pH6.6 was prepared by adding 62.50ml 0.2M monobasic stock and 37.50ml 0.2M dibasic stock in 200ml distilled water. The pH of the solution was calibrated to pH6.6 by using a pH meter in the laboratory.

3. β -carotene bleaching assay

0.2mg/ml β -carotene solution

0.2 mg/ml of β -carotene solution was prepared by dissolving 0.002 mg of β -carotene in 10 ml chloroform. The solution was kept in a centrifuge tube and wrapped in aluminium foil to avoid any unwanted degradation to the solution by light.

Tween 80

Tween 80 (polyoxyethylene sorbitan monolaurate)

Linoleic acid

Linoleic acid is obtained from Sigma. Best kept in ice throughout experiment and kept in -20°C when not in use.

4. Metal Chelating Assay

Preparation of EDTA (Ethylenediaminetetraacetic acid)

1 g of EDTA is dissolved in 4 ml distilled water. The pH is adjusted to pH 8 by using sodium hydroxide (NaOH) while stirring. EDTA will start dissolve completely once the pH has reach pH 8. Once the EDTA dissolved, the total volume is made up to 10 ml with distilled water.

Preparation of Ferrozine (3-(2-Pyridyl)-5,6-diphenyl-1,2,4-triazine-*p,p'*-disulfonic acid monosodium salt hydrate) (5 mM)

0.0024625 g of ferrozine was diluted with deionized distilled water.

Preparation of ferric chloride (FeCl_3) (2 mM)

0.039762 g of ferric chloride crystal which is light green in colour is diluted with 100 ml of deionized distilled water. This mixture must be left standing for some time for the crystals to be completely diluted.

Preparation of deionized water

Deionized water (18 megaohm-cm) was prepared using machine. Deionized water is used to avoid contaminations by ions outside of the reaction system that might affect the results of the experiments.

5. Superoxide dismutase (SOD) activity assay

Preparation of working solutions

WST working solution

This is prepared by diluting 1 ml of WST solution with 19 ml of Buffer Solution. This solution has the ability to be stable for almost 2 months at the temperature 4°C.

Enzyme Working Solution

The Enzyme Solution tube was initially centrifuged for 5 seconds. The Enzyme Solution (15 µl) was then pipetted and mixed with 2.5 ml of dilution buffer. This solution was stable for 3 weeks at 4°C.

6. Total Phenolic Content (Folin-Ciocalteu Assay)

Preparation of Gallic Acid Solution

0.5 g of dry gallic acid, which is also known as 3,4,5-trihydroxybenzoic acid is diluted with 10ml of absolute ethanol. The mixture is then made into 100ml with distilled water. This preparation can be kept in refrigerator for 2 weeks.

Preparation of Saturated Sodium Carbonate Solution (~35%)

200 g of anhydrous sodium carbonate is added into 800 ml of water. It is then brought into boiling and then left to cool thereafter. After the solution is cooled, a few crystals of sodium carbonate were added and it is left standing at room temperature for 24 hours. After 24 hours, the thick sodium carbonate solution is filtered and water is added to make the whole solution to 1 liter.

Folin-Ciocalteu Phenol Reagent (Sigma)

It contains phosphomolybdate and phosphotungstate. Commonly used for colorimetric assay to determine phenolic and polyphenolic antioxidants.

7. Thiobarbituric Acid Reactive Species (TBARS) Assay

Egg yolk suspension

A fresh egg was purchased and rinsed. The egg was broken and egg white is discarded. Only the egg yolk was taken and used for the experiment. The egg yolk was weighed and the weight of the egg yolk noted. Dilution with PBS was conducted at the ratio 1:1. The diluted egg yolk was then mixed well using a magnetic stirrer. This egg yolk was then diluted again 40x using PBS (pH 7.4) to obtain yolk suspension.

Method of egg yolk suspension preparation

Let say the weight of egg yolk is 13.4 g

The egg yolk would be diluted with PBS (pH 7.4) at 1:1 ratio; 13.4 g was added with 13.4 ml of PBS. This is then stirred using a magnetic stirrer.

To obtain egg suspension, the diluted egg yolk was then diluted again 40x (1 ml of egg yolk added into 39 ml of PBS) using PBS. In this experiment, 3 ml of the diluted egg yolk is added to 117 ml of PBS.

0.8% 2-Thiobarbituric acid

2-Thiobarbituric acid (0.8 g) was dissolved in 100 ml distilled water. The thiobarbituric acid which was in powder form was mashed finely to enable fast dilution. Dilution of this material was quite slow as it had to be placed at 50°C for 1 hour or more until the 2-thiobarbituric acid fully dissolved.

20% Trichloroacetic acid

Trichloroacetic acid crystals (20.0 g) was dissolved in 100 ml distilled water.

24 mM FeSO₄

Light green colored FeSO₄ crystals (0.334 g) was dissolved in 50 ml of distilled water.

8. Tyrosinase Inhibitory Assay**Mushroom tyrosinase**

The mushroom tyrosinase purchased was in the concentration 3933 units/mg. Hence, 1 mg of mushroom tyrosinase was diluted with 3.933 ml phosphate buffer (pH 6.8), to obtain mushroom tyrosinase stock with a concentration 1000 units/ml.

0.1 M Phosphate buffer (pH 6.8)

0.2M monobasic stock

13.9 g of sodium phosphate anhydrous monobasic of 15.6 sodium phosphate monobasic dehydrate is added with 500 ml distilled water

0.2M dibasic stock

53.65 g sodium phosphate dibasic heptahydrate is diluted with 1L distilled water or 28.4 g of anhydrous sodium phosphate dibasic with 1L distilled water.

0.1M phosphate buffer, pH 6.8

153.0 ml monobasic stock is added with 147.0 ml and 300 ml of distilled water is added. The pH is adjusted to 6.8 using a pH meter.

L-Tyrosine

L-tyrosine which acts as substrate is prepared at a concentration of 2.0mM. L-tyrosine powder (Sigma) (0.036 g) was diluted with 100 ml of 0.1M phosphate buffer solution (pH 6.8).

L-DOPA

L-DOPA which acted also as substrate was prepared at concentration of 12.0 mM. 0.2366 g of L-DOPA in powder form purchased from Sigma was diluted with 100 ml of 0.1 M phosphate buffer solution (pH 6.8).

Cytotoxicity Studies

Neutral Red Cytotoxicity Assay

Neutral Red Stock Solution

0.4g of Neutral Red (ICN, USA) was dissolved in 100 ml distilled water. The solution was kept at 4°C.

Neutral Red Medium

The Neutral Red stock solution (4mg/ml) was diluted (250µl) in treatment culture medium (20ml) to give a final concentration of 50µg/ml. The solution is then wrapped with aluminium foil and then centrifuged at 1000 rpm for 10 minutes before use to remove any fine, needle like precipitate of dye crystals.

Neutral Red Washing Solution

1.0g of calcium chloride (CaCl₂) Sigma was dissolved in 500µl formaldehyde (Sigma) and 99.5 ml distilled water. Solution was kept at 4°C.

Neutral Red Resorb Solution

1 ml of glacial acetic acid (BDH, AnalaR) was dissolved in 50 ml of absolute ethanol and 49 ml of distilled water.

Solution was kept at 4°C.

Basic (McCOY'S 5A, RPMI, MEM and M199) Medium

Medium was prepared by dissolved 11.9 g of (McCOY'S 5A, RPMI, MEM and M199) powder, 0.5206 g HEPES sodium salt (2mM/ L) and 2.0 g of sodium bicarbonate (NaHCO₃) in 1000 ml of distilled water. Then pH of the medium was calibrated to pH 7.4. The medium was then filter sterilized through a 0.22 µm filter membrane into sterile bottles and kept at 4°C.

10% supplemented medium (for cell maintenance)

10% of Supplemented Medium was prepared by using 90 ml of basic medium, supplemented with 10 ml inactivated Foetal Bovine Serum (FBS), 2 ml penicillin or streptomycin and 1 ml of Amphostat B (fungizone). The medium were filter sterilized using a 0.22 µm filter membrane. The colours of the medium were reddish orange and were kept at 4°C for up to 2 weeks.

20% Supplemented Medium (for cell reviving)

20% of Supplemented Medium were prepared by using 50 ml of 10% Supplemented Medium added with 5 ml inactivated Foetal Bovine Serum (FBS). The medium were then filter sterilized using a 0.22 µm filter membrane. The colours of the medium were reddish orange and were kept at 4°C for up to 2 weeks.

Phosphate Buffered Saline (PBS)

Preparation was done using 1.52 g of sodium phosphate anhydrous (Na₂HPO₄, BHD), 0.58 g of potassium dihydrogen orthophosphate (KH₂PO₄, BHD) and 8.5 g of sodium chloride (NaCl, Merck, Germany) that were dissolved in distilled water and the volume was made

up to 1 liter. The pH of the buffer was adjusted to 7.2. The buffer was then filtered using a 0.02 µm filter and stored at room temperature.

Phosphate Buffered Saline (PBS)

The Phosphate buffered saline (PBS) was prepared by dissolving 1.52g sodium phosphate anhydrous (NaHPO_4 , Merck, Germany), 0.58g potassium dihydrogen orthophosphate (KH_2PO_4 , Merck, Germany) and 8.5g sodium chloride (NaCl , BDH, AnalaR) in distilled water and the volume was made up to 1000ml. The pH of the buffer was adjusted to 7.2 by adding base or acid and monitored using pH meter. The buffer was then filtered by using a 0.22µm filter membrane and stored at room temperature.

Trypsin - EDTA

Trypsin - EDTA solution was prepared by dissolving 0.25g trypsin (Amresco, USA) and 0.03g EDTA (Sigma, USA) in 100ml of distilled water. The solution was sterilized by filtration using a 0.22µm filter membrane and stored at -20°C.

0.4% Tryphan Blue

0.4% Tryphan Blue solution was prepared by dissolving 0.2g tryphan blue in 50ml distilled water.

APPENDIX 3: Preparation of Apparatus

Washing of Apparatus

Glass and pipettes were soaked overnight in a diluted solution of 7X detergent (Flow Laboratories). They were rinsed thoroughly in tap water and followed by distilled water before drying in an oven (69°C).

Sterilization

Pipettes were plugged with cotton wool and placed in metal canisters. They were sterilized in hot air oven at 180°C for 2 hours. Bottles, pipettor tips and filter units were autoclaved at 15 psi at 121°C for 20 minutes.